

<p><b><u>Nitroglycerin</u></b></p> <p><b>Indications:</b></p> <ul style="list-style-type: none"> <li>• Angina Pectoris</li> <li>• Hypertension (diastolic BP &gt; 120)</li> <li>• Pulmonary Edema</li> </ul> <p><b>Pearls:</b></p> <ul style="list-style-type: none"> <li>• May cause hypotension, pulsating headache, flushing</li> <li>• Do not give if BP is &lt; 90 systolic</li> <li>• Use in hypertension requires Medical Control Physician order</li> </ul>	<p><b>Adult:</b></p> <ul style="list-style-type: none"> <li>• 1 spray or 0.4 mg tablet SL</li> <li>• Repeat every 5 minutes until pain free or max. 3 doses</li> </ul> <p><b>Pediatric:</b></p> <ul style="list-style-type: none"> <li>• N/A</li> </ul>
<p><b><u>Oxygen</u></b></p> <p><b>Indications:</b></p> <ul style="list-style-type: none"> <li>• Useful in airway, chest pain, and respiratory distress. Required for preoxygenation prior to intubation</li> </ul> <p><b>Pearls:</b></p> <ul style="list-style-type: none"> <li>• As indicated in Treatment protocols</li> </ul>	<p><b>Adult:</b></p> <ul style="list-style-type: none"> <li>• 1-4 LPM via nasal cannula</li> <li>• 6-15 LPM via NRB mask</li> <li>• 15 liters via BVM</li> </ul> <p><b>Pediatric:</b></p>
<p><b><u>Pralidoxime (2-PAM)</u></b></p> <p>(See WMD protocol)</p> <p><b>Indications:</b></p> <ul style="list-style-type: none"> <li>• Antidote for Nerve Agents or Organophosphate</li> </ul> <p><b>Pearls:</b></p> <ul style="list-style-type: none"> <li>• Administered with Atropine</li> <li>• This drug is a non-stock item and may be used upon availability</li> </ul>	<p><b>Adult:</b></p> <ul style="list-style-type: none"> <li>• Major symptoms - 600 mg IM x 3</li> <li>• Minor symptoms - 15-25 mg/kg IV over 30 minutes</li> </ul> <p><b>Pediatric:</b></p> <ul style="list-style-type: none"> <li>• Major symptoms: <ul style="list-style-type: none"> <li>0- 7 years old - 600 mg IM</li> <li>8-14 years old - 600 mg x 2</li> <li>&gt; 15 years old - 600 mg x 3</li> </ul> </li> <li>• Minor Symptoms: 15-25 mg/kg IV over 30 minutes.</li> </ul>
<p><b><u>Promethazine</u></b></p> <p><b>Indications:</b></p> <ul style="list-style-type: none"> <li>• Motion sickness, nausea, vomiting</li> </ul> <p><b>Pearls:</b></p> <ul style="list-style-type: none"> <li>• May cause drowsiness and hypotension, especially in the elderly</li> <li>• Dilute IV phenergan in 10 ml of Normal Saline Solution</li> </ul>	<p><b>Adult:</b></p> <ul style="list-style-type: none"> <li>• 12.5 mg IV over 2 min. or deep IM</li> <li>• (&gt;60 yrs old) 6.25 mg IV over 2 min. or deep IM</li> </ul> <p><b>Pediatric:</b></p> <ul style="list-style-type: none"> <li>• 0.25-0.5 mg/kg IV over 2 min. or deep IM</li> </ul>